

Aspects of hadronic J/ψ production *

K. Sridhar[†]

*Theory Group, Tata Institute of Fundamental Research,
Homi Bhabha Road, Bombay 400 005, India.*

ABSTRACT

In this talk, I start with a brief introduction to Non-Relativistic QCD (NRQCD) and its applications to quarkonium physics. This theory has provided a consistent framework for the physics of quarkonia, in particular, the colour-octet Fock components predicted by NRQCD have important implications for the phenomenology of charmonium production in experiments. We discuss the applications of NRQCD to J/ψ production at Tevatron and the tests of the theory in other experiments. Finally we discuss the production of the 1P_1 charmonium state, h_c , at large- p_T at the Tevatron. The observation of h_c is interesting, since its existence has yet to be experimentally confirmed. Moreover, this rate is a *prediction* of NRQCD, and the observation of the h_c at the Tevatron can, therefore, be used as a test of NRQCD.

*Presented at the Pacific Particle Physics Phenomenology (P^4 '97) Conference at the Seoul National University, S. Korea, October 31 - November 2, 1997.

[†]sridhar@theory.tifr.res.in

J/ψ production has traditionally been described in terms of the colour-singlet model [1, 2], which, in spite of its reasonable success with its predictions at lower energies, fails completely in explaining the data on large- p_T J/ψ production at the Tevatron [3]. This has led to a major revision of the theory of quarkonium formation. In the colour-singlet model, the relative velocity, v between the charm quarks is neglected. However, for charmonia $v^2 \sim 0.3$, and is not negligible. It is possible to include the effects coming from higher orders in a perturbation series in v in non-relativistic QCD (NRQCD) [4, 5].

In this picture, the physical charmonium state can be expanded in terms of its Fock-components in a perturbation series in v , and the $c\bar{c}$ states appear in either colour-singlet or colour-octet configurations in this series. The colour-octet $c\bar{c}$ state makes a transition into a physical state by the emission of one or more soft gluons, dominantly *via* $E1$ or $M1$ transitions with higher multipoles being suppressed by powers of v . The cross-section for the production of a meson H then takes on the following factorised form:

$$\sigma(H) = \sum_{n=\{\alpha,S,L,J\}} \frac{F_n}{m^{d_n-4}} \langle \mathcal{O}_\alpha^H(^{2S+1}L_J) \rangle \quad (1)$$

where F_n 's are the short-distance coefficients and $\mathcal{O}_n$ are local 4-fermion operators, of naive dimension d_n , describing the long-distance physics.

In fact, the importance of the colour-octet components was first noted [6] in the case of P -wave charmonium decays, and even in the production case the importance of these components was first seen [7] in the production of P -state charmonia at the Tevatron. The surprise was that even for the production of S -states such as the J/ψ or ψ' , where the colour singlet components give the leading contribution in v , the inclusion of sub-leading octet states was seen to be necessary for phenomenological reasons [8]. While the inclusion of the colour-octet components seem to be necessiated by the Tevatron charmonium data, the normalisation of these data cannot be predicted because the the long-distance matrix elements are not calculable. The data allow a linear combination of octet matrix-elements to be fixed [9, 10], and much effort has made recently to understand the implications of these colour-octet channels for J/ψ production in other experiments. We discuss some of these below.

1. The prediction [11, 12, 13] for prompt J/ψ production at LEP in the colour-singlet model is of the order of 3×10^{-5} , which is almost an order of magnitude below the experimental number for the branching fraction obtained from LEP [14]. Recently, the colour-octet contributions to J/ψ production in this channel have been studied [15, 16] and it is found that the inclusion of the colour-octet contributions in the fragmentation functions results in a predictions for the branching ratio which is 1.4×10^{-4} which is compatible with the measured values of the branching fraction from LEP [14].
2. The production of J/ψ in low energy e^+e^- machines can also provide a stringent test of the colour-octet mechanism [17]. In this case, the colour-octet contributions dominate near the upper endpoint of the J/ψ energy spectrum, and the signature for the colour-octet process is a dramatic change in the angular distribution of the J/ψ near the endpoint.
3. One striking prediction of the colour-octet fragmentation process both for $p\bar{p}$ colliders and for J/ψ production at the Z -peak, is that the J/ψ coming from the process $g \rightarrow J/\psi X$ is produced in a transversely polarised state [18]. For the colour-octet $c\bar{c}$ production, this is predicted to be a 100% transverse polarisation, and heavy-quark spin symmetry will then ensure that non-perturbative effects which convert the $c\bar{c}$ to a J/ψ will change this polarisation only very mildly. This spin-alignment can, therefore, be used as a test of colour-octet fragmentation.

4. The colour-octet components are found [19] to dominate the production processes in fixed-target pp and πp experiments. Using the colour-octet matrix elements extracted from elastic photoproduction data it is possible to get a very good description of the $\sqrt{s}$ -dependence and also the x_F and rapidity distributions.
5. The associated production of a $J/\psi + \gamma$ is also a crucial test of the colour-octet components [20] and also of the fragmentation picture [21]. Similar tests can be conceived of with double J/ψ production at the Tevatron [22].
6. J/ψ and ψ' production in pp collisions at centre-of-mass energies of 14 TeV at the LHC also provides a crucial test of colour-octet fragmentation [23].
7. The fragmentation process has also definite predictions for the large- p_T inelastic J/ψ photoproduction at HERA [24]. At HERA, the colour-singlet component dominates inelastic J/ψ production but at large- z ($z > 0.7$) the colour-octet components start growing rapidly. Such a rapid growth is clearly in conflict with experiment [25]. The applicability of NRQCD factorisation at these large values of z has been questioned [26], and more work needs to be done before the disagreement between theory and the HERA inelastic data can be interpreted as a failure of NRQCD.

One important feature of the NRQCD Lagrangian is that it shows an approximate heavy-quark symmetry, which is valid to $O(v^2) \sim 0.3$. The implication of this symmetry is that the nonperturbative parameters have a weak dependence on the magnetic quantum number. We show that the production cross-section of the 1P_1 charmonium state, h_c , can be *predicted* in NRQCD, because of the heavy-quark symmetry. The Tevatron data on χ_c production fixes [10] the colour-octet matrix element which specifies the transition of a 3S_1 octet state into a 3P_J state. We would expect from heavy-quark spin symmetry of the NRQCD Lagrangian that the matrix-element for $^1S_0^{[8]} \rightarrow h_c$ should be of the same order as that for $^3S_1^{[8]} \rightarrow ^3P_1$. The production of the 1P_1 charmonium state, h_c , is, therefore, a crucial prediction of NRQCD.

The production of the h_c is interesting in its own right : charmonium spectroscopy [27] predicts this state to exist at the centre-of-gravity of the $\chi_c(^3P_J)$ states. While the E760 collaboration at the Fermilab has reported [28] the first observation of this resonance its existence needs further confirmation. In the following, we report on the calculations for h_c production at the Tevatron presented in Ref. [29].

Being a P -state, the leading colour-singlet contribution is already at $O(v^2)$, and at the same order we have the octet production of the 1P_1 state through an intermediate 1S_0 state. The octet channel is, therefore, expected to be very important for the production of this state. Once produced, the h_c can be detected by its decay into a $J/\psi + \pi^0$. The subprocesses that we are interested in are the following:

$$\begin{aligned}
g + g &\rightarrow ^1P_1^{[1]} + g, \\
g + g &\rightarrow ^1S_0^{[8]} + g, \\
q(\bar{q}) + g &\rightarrow ^1S_0^{[8]} + q(\bar{q}), \\
q + \bar{q} &\rightarrow ^1S_0^{[8]} + g.
\end{aligned} \tag{2}$$

The $^1S_0^{[8]} \rightarrow h_c$ is mediated by a gluon emission in a $E1$ transition.

The cross-section $d\sigma/dp_T$ for h_c production at the Tevatron energy ($\sqrt{s} = 1.8$ TeV) presented in Ref. [29] are shown in Fig. 1. For 20 pb^{-1} total luminosity, for p_T integrated between 5 and 20 GeV we expect of the order of 650 events in the $J/\psi + \pi$ channel. Of these, the contribution from the colour-singlet channel is a little more than 40, while the octet channel gives

more than 600 events. The colour-octet dominance is more pronounced at large- p_T . Recent results on J/ψ production from CDF are based on a total luminosity of 110 pb^{-1} . For this sample, more than 3000 events can be expected to come from the decay of the h_c into a J/ψ and a π . With this large event rate, the h_c should certainly be observable if the π^0 coming from its decay can be reconstructed efficiently.

In conclusion, we find that a reasonably large rate for the production of the 1P_1 is expected at the Tevatron, with a dominant contribution from the colour-octet production channel. Heavy-quark symmetry relations allow us to infer the size of the non-perturbative matrix-elements and to predict the rate for h_c production and its subsequent decay into a J/ψ and a π . By looking at J/ψ events associated with a soft pion, it should be possible to pin down the elusive 1P_1 resonance at the Tevatron, given the large number of J/ψ events already available. We believe that this is a firm prediction of the NRQCD framework and may be a useful way of distinguishing this from other models of quarkonium formation.




Figure 1: $d\sigma/dp_T$ (in nb/GeV) for h_c production (after folding in with $\text{Br}(h_c \rightarrow J/\psi + \pi)=0.5$) at 1.8 TeV c.m. energy with $-0.6 \leq y \leq 0.6$.

References

- [1] E.L. Berger and D. Jones, *Phys. Rev.* **D 23** (1981) 1521.
- [2] R. Baier and R. Rückl, *Z. Phys.* **C 19** (1983) 251.
- [3] F. Abe et al., *Phys. Rev. Lett.* **69** (1992) 3704; *Phys. Rev. Lett.* **71** (1993) 2537; R. Demina, FERMILAB-CONF-96-201-E, Presented at the 11th Topical Workshop on Proton-Antiproton Collider Physics (PBARP 96), Padua, Italy, 26th May – 1st June 1996.
- [4] W.E. Caswell and G.P. Lepage, *Phys. Lett.*, **B 167** (1986) 437.
- [5] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev.* **D 51** (1995) 1125.
- [6] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev.* **D 46** (1992) R1914.

- [7] E. Braaten, M.A. Doncheski, S. Fleming and M. Mangano, *Phys. Lett. B* **333** (1994) 548; D.P. Roy and K. Sridhar, *Phys. Lett. B* **339** (1994) 141; M. Cacciari and M. Greco, *Phys. Rev. Lett.* **73** (1994) 1586.
- [8] E. Braaten and S. Fleming, *Phys. Rev. Lett.* **74** (1995) 3327.
- [9] M. Cacciari, M. Greco, M. Mangano and A. Petrelli, *Phys. Lett. B* **356** (1995) 553.
- [10] P. Cho and A.K. Leibovich, *Phys. Rev.*, **D 53** (1996) 150; *Phys. Rev.*, **D 53** (1996) 6203.
- [11] V. Barger, K. Cheung and W.-Y. Keung, *Phys. Rev. D* **41** (1990) 1541.
- [12] W.-Y. Keung, *Phys. Rev. D* **23** (1981) 2072; J.H. Kühn and H. Schneider, *Phys. Rev. D* **24** (1981) 2996.
- [13] E. Braaten, K. Cheung and T.C. Yuan, *Phys. Rev. D* **48** (1993) 4230.
- [14] O. Adriani et al. (L3 collaboration), *Phys. Lett. B* **288** (1992) 412; P. Abreu et al. (DELPHI collaboration), *Phys. Lett. B* **341** (1994) 109; G. Alexander et al. (OPAL collaboration), CERN Preprint CERN-PPE-95-153.
- [15] K. Cheung, W.-Y. Keung and T.C. Yuan, *Phys. Rev. Lett.* **76** (1996) 877; S. Baek, P. Ko, J. Lee and H.S. Song, *Phys. Lett. B* **389** (1996) 609.
- [16] P. Cho, *Phys. Lett. B* **368** (1996) 171.
- [17] E. Braaten and Y. Chen, *Phys. Rev. Lett.* **76** (1996) 730.
- [18] P. Cho and M. Wise, *Phys. Lett. B* **346** (1995) 129; S. Baek, P. Ko, J. Lee and H.S. Song, *Phys. Rev. D* **55** (1997) 6839; M. Beneke and I. Rothstein, *Phys. Lett. B* **372** (1996) 157; M. Beneke and M. Krämer, *Phys. Rev. D* **55** (1997) 5269.
- [19] S. Gupta and K. Sridhar, *Phys. Rev. D* **54** (1996) 5545; M. Beneke and I. Rothstein, *Phys. Rev. D* **54** (1996) 2005; S. Gupta and K. Sridhar, *Phys. Rev. D* **55** (1997) 2650.
- [20] C.S. Kim, J. Lee and H.S. Song, KEK Preprint KEK-TH-474, hep-ph/9610294.
- [21] D.P. Roy and K. Sridhar, *Phys. Lett. B* **341** (1995) 413.
- [22] V. Barger, S. Fleming and R.J.N. Phillips, *Phys. Lett. B* **371** (1996) 111.
- [23] K. Sridhar, *Mod. Phys. Lett. A* **11** (1996) 1555.
- [24] R.M. Godbole, D.P. Roy and K. Sridhar, *Phys. Lett. B* **373** (1996) 328.
- [25] N. Cacciari and M. Krämer, *Phys. Rev. Lett.* **76** (1996) 4128; P. Ko, J. Lee, H.S. Song, *Phys. Rev. D* **54** (1996) 4312.
- [26] M. Beneke, I.Z. Rothstein, M.B. Wise, *Phys. Lett. B* **408** (1997) 373.
- [27] Y.-P. Kuang, S.F. Tuan and T.-M. Yan, *Phys. Rev. D* **37** (1988) 1210.
- [28] T.A. Armstrong et al., *Phys. Rev. Lett.* **69** (1992) 2337.
- [29] K. Sridhar, *Phys. Rev. Lett.* **77** (1996) 4880.